

Summary

Week 17/2017 (24 – 30 April 2017)

- Influenza activity across the region decreased further and all 40 reporting countries reported low influenza activity.
- The overall proportion of sentinel specimens testing positive for influenza viruses returned to the epidemic threshold value (10%).
- Type B viruses represented 95% of sentinel detections. However, the overall number of type B virus detections remained low.

Season overview

- After an earlier start than usual (week 46/2016), the influenza season is considered to be over in the great majority of countries in the Region, with influenza activity at inter-seasonal levels in all reporting countries and with the proportion of sentinel detections returning to the epidemic threshold value (10%).
- From week 40/2016 through week 10/2017, influenza A viruses predominated, accounting for 90% of all sentinel detections, and 99% of those subtyped were A(H3N2). Since week 11/2017, influenza B viruses have predominated, although absolute numbers of type B detections have remained low.
- Confirmed cases of influenza type A virus infection reported from hospitals have predominantly been in adults aged 65 years or older.
- Significant excess all-cause mortality has been observed in people aged 15–64 years, and markedly so in people aged 65 years or older, in the majority of the 20 reporting countries or regions. This is commonly seen when the predominant viruses circulating are A(H3N2).
- Two-thirds of the A(H3N2) viruses genetically characterized belong to subclade 3C.2a1, but remain antigenically similar to the clade 3C.2a vaccine virus, as described in the [WHO recommendations for vaccine composition for the northern hemisphere 2017–18](#). [See also the WHO CC London February 2017 report](#).
- Vaccine effectiveness estimates for all age groups against A(H3N2) illness suggest moderate effectiveness in [Canada](#) (42%), the [US](#) (43%) and in [Europe](#) (38%).
- Of the viruses tested so far this season, one A(H3N2) virus has shown reduced susceptibility to oseltamivir and another A(H3N2) virus has shown reduced susceptibility to zanamivir.
- The developments during the season have been consistent with the conclusions of the ECDC [risk assessment](#) on seasonal influenza, [updated](#) on 25 January 2017, which suggested increased severe outcomes in the elderly due to the high prevalence of A(H3N2) viruses, resulting in pressure on some health care systems.

Primary care data

Influenza activity

All 40 countries reporting on influenza activity for week 17/2017 reported low intensity (Fig. 1).

However, of the 40 countries reporting on geographic spread of influenza, 1 reported widespread, 2 regional and 22 local or sporadic influenza activity, indicating that influenza viruses are still circulating; 15 countries reported no influenza activity (Fig. 2).

The proportion of influenza virus detections among sentinel specimens has returned to the epidemic threshold value (10%) with 11 of 34 countries reporting dominance of influenza B viruses.

Maps of qualitative indicators in the European Region

Fig. 1. Intensity in the European Region, week 17/2017





Fig. 2. Geographic spread in the European Region, week 17/2017



For interactive maps of influenza intensity and geographic spread, please see the Flu News Europe [website](#).

Viruses detected in sentinel-source specimens (ILI and ARI)

For week 17/2017, 41 (10%) of 421 sentinel specimens tested positive for influenza viruses (Table 1). Of these, 95% were type B and 5% type A viruses. The proportion of type B viruses commonly increases in the second half of an influenza season. Of 10 countries across the region that each tested at least 10 sentinel specimens, 4 countries reported a proportion of influenza virus detections of 10% or above (Armenia, Republic of Moldova, Turkey and the United Kingdom (Scotland)). The 2 type A viruses were sub-typed as A(H3N2). Of 8 influenza B viruses ascribed to a lineage, 7 were B/Victoria and 1 was B/Yamagata.

Since week 40/2016, of all typed viruses, 90% were type A, with 99% of those subtyped being A(H3N2) (Fig. 3, Table 1). Of the 771 influenza B viruses that have been ascribed to a lineage since week 40/2016, 431 (56%) were of the B/Yamagata lineage and 340 (44%) were of the B/Victoria lineage.

Fig. 3. Influenza virus detections in sentinel-source specimens by type and subtype, by week




Table 1. Influenza virus detections in sentinel-source specimens by type and subtype, week 17/2017 and cumulatively

Virus type and subtype	Current Week		Season 2016-2017	
	Number	% ^a	Number	% ^a
Influenza A	2	5	16 235	90
A(H1N1)pdm09	0	0	186	1
A(H3N2)	2	100	13 567	99
A not subtyped	0	-	2 482	-
Influenza B	39	95	1 834	10
B/Victoria lineage	7	90	340	44
B/Yamagata lineage	1	10	431	56
Unknown lineage	31	-	1 063	-
Total detections / Total tested	41 / 421	10	18 069 / 49 931	36

^aFor influenza type percentage calculations, the denominator is total detections; for subtype and lineage, it is total influenza A subtyped and total influenza B lineage determined, respectively; for total detections, it is total tested.

Severity

8 of the 10 countries that reported data on sentinel surveillance of severe acute respiratory infections (SARI) reported 516 SARI cases for week 17/2017. Among these cases, 131 respiratory specimens were collected, 13 (10%) of which tested positive for influenza viruses in 4 countries (Armenia, Georgia, Republic of Moldova and the Russian Federation).

Since week 40/2016, 16 countries have reported 38 328 SARI cases. Of these 10 495 were tested for influenza viruses, 3 540 (34%) of which were positive: 2 706 (76%) were type A

and 834 (24%) type B viruses. Of the influenza A viruses, 2 493 (92%) were A(H3N2), 7 (<1%) were A(H1N1)pdm09 and 206 (8%) were not subtyped.

For week 17/2017, the 9 countries that conduct surveillance of hospitalized laboratory-confirmed influenza cases reported no cases.


Since week 40/2016, the 9 countries reported 3 654 cases that have been admitted to ICU; 3 561 (97%) were infected with influenza type A viruses (2 117 - unsubtyped, 1 317 - A(H3N2) and 127 - A(H1N1)pdm09) and 93 with type B viruses.

Since week 40/2016, 5 countries have reported 3 741 laboratory-confirmed influenza cases admitted to non-ICU wards; 3 693 (99%) were infected with influenza type A viruses (2 063 - unsubtyped, 1 623 - A(H3N2), 7 - A(H1N1)pdm09), and 48 were infected with type B influenza viruses.

Since the start of the season, information on patient age and influenza virus (sub)types was available for 2 598 cases admitted to ICU; the majority of cases (64%; n=1 665) were aged ≥65 years, 791 (31%) were aged 15–64 years and 142 (5%) were aged under 15 years (**Error! Not a valid bookmark self-reference.**).

In total, 931 deaths have been reported, 532 from ICUs and 399 from non-ICU wards, with 758 (81%) of all deaths occurring in patients aged 65 years or older. Of all fatal cases, 916 (99%) were due to influenza A with 452 (99%) of those subtyped being A(H3N2) viruses.

Fig. 4. Distribution of virus (sub)types in influenza-confirmed cases admitted to ICU by age-group, cumulatively, during weeks 40/2016-17/2017



Mortality monitoring

Data from 20 countries or regions reporting to the [Euromomo](#) project were received for week 17/2017 and included in the pooled analyses of excess all-cause mortality.

The majority of participating European countries experienced a [marked excess](#) in all-cause mortality between the beginning of January 2017 and the end of February 2017, in particular among the elderly (those aged 65 years and above). Mortality levels have since decreased to expected levels. This season's excess mortality coincided with circulation of influenza A(H3N2) viruses, which usually leads to increased mortality among the elderly.


Virus characteristics

Viruses detected in non-sentinel-source specimens

For week 17/2017, 705 specimens from non-sentinel sources (such as hospitals, schools, non-sentinel primary care facilities, nursing homes and other institutions) tested positive for influenza viruses (Fig. 5, Table 2).

Of these, 17% were type A (with all subtyped viruses being A(H3N2)), and 83% type B. The increase in proportion of type B viruses corresponds to the sentinel detection data, however the number of influenza B viruses detected remained relatively low and similar to that seen in recent weeks.

Fig. 5. Influenza virus detections in non-sentinel-source specimens by type and subtype, by week



While subtype and lineage were not determined for the majority of influenza viruses since week 40/2016, cumulative distributions of types and type A subtypes similar to those among sentinel detections have been observed: of all typed viruses, 88% were type A, with 99% of those subtyped being A(H3N2). Of 1 380 influenza type B viruses ascribed to a lineage, 77%

were B/Yamagata lineage and 23% were B/Victoria lineage (Table 2), which differs from sentinel detections where B/Victoria lineage and B/Yamagata lineage viruses have been more evenly distributed this season. The difference is mainly driven by the proportion of influenza B lineage detections in sentinel specimens in Latvia, Norway and Slovenia (B/Yamagata lineage predominant).

Table 2. Influenza viruses detected in non-sentinel-source specimens, by virus (sub)type, week 17/2017 and cumulatively

Virus type and subtype	Current Week		Season 2016-2017	
	Number	% ^a	Number	% ^a
Influenza A	123	17	109 698	88
A(H1N1)pdm09	0	0	359	1
A(H3N2)	30	100	39 211	99
A not subtyped	93	-	70 128	-
Influenza B	582	83	14 541	12
B/Victoria lineage	0	0	322	23
B/Yamagata lineage	6	100	1 058	77
Unknown lineage	576	-	13 161	-
Total detections / Total tested	705 / 8 034	-	124 239 / 566 420	-

^aFor influenza type percentage calculations, the denominator is total detections; for subtype and lineage, it is total influenza A subtyped and total influenza B lineage determined, respectively; as not all countries have a true non-sentinel testing denominator, no percentage calculations for total tested are shown.

Genetic characterization

For specimens collected since week 40/2016, genetic characterization of 3 485 viruses has been reported (Table 3). Among 3 128 A(H3N2) viruses, 928 fell in the vaccine component clade (3C.2a) and 2 161 in the 3C.2a1 subclade defined by N171K amino acid substitution, often with N121K, in the haemagglutinin. Viruses in these two clades have been antigenically similar, but both clades are evolving rapidly with the emergence of several virus clusters defined by additional amino acid substitutions in the haemagglutinin, thereby requiring continued monitoring of antigenic characteristics. See also the [WHO CC London February 2017 report](#).

Table 3. Viruses attributed to genetic groups, cumulative for weeks 40/2016–17/2017

Phylogenetic group	Number of viruses
A(H1N1)pdm09 A/Michigan/45/2015 (subgroup 6B.1) ^{b, c}	29
A(H1N1)pdm09 A/South Africa/3626/2013 (subgroup 6B)	5
A(H3N2) A/Bolzano/7/2016 (subgroup 3C.2a1)	2161
A(H3N2) A/Hong Kong/4801/2014 (subgroup 3C.2a) ^{a, b, c}	928
A(H3N2) A/Samara/73/2013 (subgroup 3C.3)	1
A(H3N2) A/Switzerland/9715293/2013 subgroup (3C.3a)	31
A(H3N2) A/Stockholm/28/2014 (subgroup 3C.3b)	1
A(H3N2), subgroup not listed	6
B/Brisbane/60/2008 (Victoria lineage clade 1A) ^{a, b, c}	73
B/Phuket/3073/2013 (Yamagata lineage clade 3) ^d	240

^a Vaccine component for Northern Hemisphere 2016–2017 season

^b Vaccine component for Southern Hemisphere 2017 season

^c Vaccine component for Northern Hemisphere 2017–2018 season

^d Vaccine component of quadrivalent vaccines for use in both Northern and Southern Hemisphere

The recommended composition of trivalent influenza vaccines for the 2016–2017 season in the [northern hemisphere](#) was for inclusion of an A/California/7/2009 (H1N1)pdm09-like virus; an A/Hong Kong/4801/2014 (H3N2)-like virus; and a B/Brisbane/60/2008-like virus (B/Victoria lineage). For quadrivalent vaccines a B/Phuket/3073/2013-like virus (B/Yamagata lineage) was recommended. On 2 March 2017, WHO announced the recommended vaccine composition for the 2017–2018 season in the [northern hemisphere](#). The recommendations matched those for the 2016–2017 season, but for the A(H1N1)pdm09 component being changed to an A/Michigan/48/2015-like virus (clade 6B.1).

Early monitoring of vaccine effectiveness (VE) in [Finland and Sweden](#) (Stockholm County) suggested levels of effectiveness in persons aged 65 years or older (32% and 28% VE, respectively) similar to estimates from annual multi-country studies covering the 2011–2012 and 2014–2015 seasons. More recent VE estimates for all age groups against A(H3N2) illness from Canada (42%), from the US (43%) and from Europe (38%) were consistent with the early estimates from Finland and Sweden.

Antiviral susceptibility testing

Neuraminidase inhibitor susceptibility has been assessed for 2 310 influenza viruses (2 137 A(H3N2), 31 A(H1N1)pdm09 and 142 type B) with collection dates since week 40/2016. All but two A(H3N2) viruses showed no phenotypic or genotypic signs for reduced inhibition (RI): one, from a specimen collected in week 01/2017, showed RI by zanamivir while the other, from a specimen collected in week 04/2017, showed RI by oseltamivir in phenotypic assays.

This weekly update was prepared by an editorial team at the European Centre for Disease Prevention and Control (Cornelia Adlhoch, Eeva Broberg, René Snacken, Pasi Penttinen) and the WHO Regional Office for Europe (Caroline Brown, Piers Mook, Dmitriy Pereyaslov and Tamara Meerhoff, Temporary Advisor to WHO). It was reviewed by country experts (AnnaSara Carnahan, Public Health Agency, Sweden; Veronica Eder, National Public Health Center, Republic of Moldova), and by experts from the network (Adam Meijer, National Institute for Public Health and the Environment (RIVM), the Netherlands; Rod Daniels and John McCauley, WHO Collaborating Centre for Reference and Research on Influenza, Francis Crick Institute, United Kingdom; Tyra Grove Krause, Statens Serum Institut and EuroMOMO network, Denmark).

Maps and commentary do not represent a statement on the legal or border status of the countries and territories shown.

All data are up to date on the day of publication. Past this date, however, published data should not be used for longitudinal comparisons, as countries retrospectively update their databases.

The WHO Regional Office for Europe is responsible for the accuracy of the Russian translation.

Suggested citation:

European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Flu News Europe, Joint ECDC–WHO weekly influenza update, week 17/2017.

Tables and figures should be referenced:

European Centre for Disease Prevention and Control/WHO Regional Office for Europe. Flu News Europe, Joint ECDC–WHO weekly influenza update, week 17/2017.

© World Health Organization 2017

© European Centre for Disease Prevention and Control 2017

Reproduction is authorized, provided the source is acknowledged.